

PRODUCT INFORMATION

Clone ID DM155 **Target** CD171

CAML1; CD171; HSAS; HSAS1; MASA; MIC5; N-**Synonyms**

CAM-L1; N-CAML1; NCAM-L1; S10; SPG1

Host Species

Description Anti-CD171 antibody(DM155); Rabbit mAb

Delivery In Stock **Uniprot ID** P32004 Rabbit IgG IgG type Clonality Monoclonal Reactivity Human

Applications ELISA; Flow Cyt

Recommended

Background

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

The protein encoded by this gene is an axonal glycoprotein belonging to the immunoglobulin supergene family. The etidodemain; consisting of several immunoglobulin-like domains and fibronectin-like repeats (type III); is linked via a single transmembrane sequence to a conserved cytoplasmic domain. This cell adhesion molecule plays an important role in nervous system

development; including neuronal migration and differentiation. Mutations in the gene cause X-

linked neurological syndromes known as CRASH (corpus callosum hypoplasia; retardation; aphasia; spastic paraplegia and hydrocephalus). Alternative splicing of this gene results in multiple transcript variants; some of which include an alternate exon that is considered to be specific to

> Email: info@dimabio.com Website: www.dimabio.com

neurons.

Research use only **Usage**






Figure 1. ELISA plate pre-coated by 1 μ g/ml (100 μ l/well) Human CD171 protein, His tagged protein ([getskuurl sku="PME100173"]) can bind Rabbit anti-CD171 monoclonal antibody(clone: DM155) in a linear range of 1-100 ng/ml.




Figure 2. CD171 protein is highly expressed on the surface of Expi293 cell membrane. Flow cytometry analysis with Anti-CD171 (DM155) on Expi293 cells transfected with human CD171 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram), and Isotype antibody on Expi293 transfected with irrelevant protein (Orange histogram).

Email: info@dimabio.com Website: www.dimabio.com

