

PRODUCT INFORMATION

Clone ID **DMC220** FCRL5 **Target**

Synonyms BXMAS1; CD307; CD307e; FCRH5; IRTA2; PRO820

Host Species Rabbit

Anti-FCRL5 antibody(DMC220); IgG1 Chimeric Description mAb

Delivery In Stock **Uniprot ID Q96RD9**

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human

Applications ELISA; Flow Cyt

Recommended

Background

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

This gene encodes a member of the

immunoglobulin receptor superfamily and the Fc-receptor like family. This gene and several other Fc receptor like family. This gene and several other Fc receptor-like gene members are clustered on the long arm of chromosome 1. The encoded protein is a single-pass type I membrane protein and contains 8 immunoglobulin-like C2-type domains. This gene is implicated in B cell development and lymphomogenesis. Alternatively

development and lymphomagenesis. Alternatively spliced transcript variants encoding different

> Email: info@dimabio.com Website: www.dimabio.com

isoforms have been identified.

Usage Research use only






Figure 1. Flow cytometry analysis with Anti-FCRL5 (DMC220) on Expi293 cells transfected with human FCRL5 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

