

PRODUCT INFORMATION

|                              |  |
|------------------------------|--|
| Target                       | BAFF-R   |
| Synonyms                     | BAFFR;TNFRSF13C;BAFF-R;BROMIX;CD268;CVID4;prolixin   |
| Description                  | Recombinant Human BAFF-R protein with C-terminal mouse Fc  |
| Delivery                     | In Stock   |
| Uniprot ID                   | Q96RJ3   |
| Expression Host              | HEK293   |
| Tag                          | C-Mouse Fc Tag   |
| Molecular Characterization   | BAFF-R(Ser7-Ala71) mFc(Pro99-Lys330)   |
| Molecular Weight             | The protein has a predicted molecular mass of 33.6 kDa after removal of the signal peptide.The apparent molecular mass of BAFF-R-mFc is approximately 35-55 kDa due to glycosylation.  |
| Purity                       | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.   |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.   |
| Storage & Shipping           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| Background                   | B cell-activating factor (BAFF) enhances B-cell survival in vitro and is a regulator of the peripheral B-cell population. Overexpression of Baff in mice results in mature B-cell hyperplasia and symptoms of systemic lupus erythematosus (SLE). Also, some SLE patients have increased levels of BAFF in serum. Therefore, it has been proposed that abnormally high levels of BAFF may contribute to the pathogenesis of autoimmune diseases by enhancing the survival of autoreactive B cells. The protein encoded by this gene is a receptor for BAFF and is a type III transmembrane protein containing a single extracellular cysteine-rich domain. It is thought that this receptor is the principal receptor required for BAFF-mediated mature B-cell survival. |
| Usage                        | Research use only  |






Figure 1. Human BAFF-R Protein, mFc Tag on SDS-PAGE under reducing condition.

**Human BAFF-R, mFc Tagged protein ELISA**  
0.2 µg of BAFF, hFc Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human BAFF, hFc tagged protein PME100043 can bind Human BAFF-R, mFc tagged protein (PME100044) in a linear range of 0.488-250.0 ng/ml.



**Human BAFFR, mFc Tagged protein ELISA**  
0.2 µg of Human BAFFR, mFc Tagged protein per well




Figure 3. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human BAFFR, mFc tagged protein (PME100044) can bind Anti-BAFFR Neutralizing antibody BME100045 in a linear range of 0.64-16 ng/ml.

