

PRODUCT INFORMATION

Target	CD69
Synonyms	AIM;BL-AC/P26;CLEC2C;EA1;GP32/28;MLR-3
Description	Recombinant Human CD69 Protein with N-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q07108
Expression Host	HEK293
Tag	N-Human Fc Tag
Molecular Characterization	hFc(Glu99-Ala330) CD69(Ser62-Lys199)
Molecular Weight	The protein has a predicted molecular mass of 42.1 kDa after removal of the signal peptide. The apparent molecular mass of hFc-CD69 is approximately 55-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	This gene encodes a member of the calcium dependent lectin superfamily of type II transmembrane receptors. Expression of the encoded protein is induced upon activation of T lymphocytes, and may play a role in proliferation. Furthermore, the protein may act to transmit signals in natural killer cells and platelets. [provided by RefSeq, Aug 2011]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human CD69 Protein, hFc Tag on SDS-PAGE under reducing condition.

