

## **PRODUCT INFORMATION**

IL21R **Target** 

**Synonyms** CD360;IMD56;NILR

Recombinant human IL21R protein with C-**Description** 

terminal human Fc tag

**Delivery** In Stock **Uniprot ID** Q9HBE5 **Expression Host HEK293** 

Tag C-Human Fc Tag

Molecular

**Background** 

IL21R(Cys20-Glu232) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight** 

50.7 kDa after removal of the signal peptide.The apparent molecular mass of IL2IR-hFc is

approximately 55-100 kDa due to glycosylation.

The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & Reconstitution

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a cytokine receptor for interleukin 21 (IL21). It belongs to the type I cytokine receptors, and has been shown to form a heterodimeric receptor complex with the common gamma-chain, a receptor subunit also shared by the receptors for interleukin 2, 4, 7, 9, and 15. This receptor transduces the growth promoting signal of IL21, and is important for the

proliferation and differentiation of T cells, B cells, and natural killer (NK) cells. The ligand binding of this receptor leads to the activation of multiple downstream signaling molecules, including JAK1, JAK3, STAT1, and STAT3. Knockout studies of a similar gene in mouse suggest a role for this gene in regulating immunoglobulin production. Three

> Email: info@dimabio.com Website: www.dimabio.com

alternatively spliced transcript variants have been described.

**Usage** Research use only










Figure 1. Human IL21R Protein, hFc Tag on SDS-PAGE under reducing condition.

## Human IL21R,hFc Tagged protein ELISA

0.2 µg of Human IL21R, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human IL21R Protein, hFc Tag(PME100301) can bind Anti-IL21R antibody(DMC225), IgG1 Chimeric mAb in a linear range of 3.20–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

