

PRODUCT INFORMATION

PLAUR Target

Synonyms U-PAR; uPAR; CD87

Recombinant human PLAUR protein with C-Description

terminal 6×His tag

Delivery In Stock **Uniprot ID** Q03405 **Expression Host HEK293** Tag C-6×His Tag

Molecular

Background

PLAUR(Leu23-Arg303) 6×His tag Characterization

The protein has a predicted molecular mass of

32.1 kDa after removal of the signal peptide. The apparent molecular mass of PLAUR-His is **Molecular Weight**

approximately 35-55 kDa due to glycosylation. The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes the receptor for urokinase plasminogen activator and, given its role in localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the extracellular matrix. It binds both the proprotein and mature forms of urokinase plasminogen activator and permits the activation of the

receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encoding different isoforms.

The proprotein experiences several post-translational cleavage reactions that have not yet been fully defined. [provided by RefSeq, Jul 2008]

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

Website: www.dimabio.com

Email: info@dimabio.com








Figure 1. Human PLAUR Protein, His Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

