

PRODUCT INFORMATION

Clone ID	26H2
Target	ANPEP
Synonyms	APN;CD13;GP150;LAP1;P150;PEPN
Host Species	Rabbit
Description	Anti-ANPEP antibody(26H2), IgG1 Chimeric mAb
Delivery	In Stock
Uniprot ID	P15144
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1/100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Aminopeptidase N is located in the small-intestinal and renal microvillar membrane, and also in other plasma membranes. In the small intestine aminopeptidase N plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Its function in proximal tubular epithelial cells and other cell types is less clear. The large extracellular carboxyterminal domain contains a pentapeptide consensus sequence characteristic of members of the zinc-binding metalloproteinase superfamily. Sequence comparisons with known enzymes of this class showed that CD13 and aminopeptidase N are identical. The latter enzyme was thought to be involved in the metabolism of regulatory peptides by diverse cell types, including small intestinal and renal tubular epithelial cells, macrophages, granulocytes, and synaptic membranes from the CNS. This membrane-bound zinc metalloprotease is known to serve as a receptor for the HCoV-229E alphacoronavirus as well as other non-human coronaviruses. This gene has also been shown to promote angiogenesis, tumor growth, and metastasis and defects in this gene are associated with various types of leukemia and lymphoma. [provided by RefSeq, Apr 2020]
Usage	Research use only
Conjugate	Unconjugated



DIMA Disclaimer

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.




Figure 1. Flow cytometry analysis with 1µg/mL Anti-ANPEP (26H2) mAb on HEK293 cells transfected with human ANPEP (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).





Figure 2. Flow cytometry analysis of antigen binding of anti-human ANPEP mAb(DMC101149).
(A) DMC101149 does not bind to CHO-S cells that do not express ANPEP.
(B) A clear peak shift of DMC101149 was seen compared to the control when incubated with ANPEP-expressing cells. Antigen binding of DMC101149 to ANPEP-expressing cells was inhibited at 5 µg/mL.

Biotechnology B7, Biolake No.666
Gaixin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Website: www.dimabio.com



DIMABIO CONFIDENTIAL

