

**PRODUCT INFORMATION**

<b>Common Name</b>	MABX-9001a
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	CD276
<b>Applications</b>	ELISA □ Flow Cyt
<b>Endotoxin</b>	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Recommended Dilutions</b>	ELISA 1:5000-10000 □ Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Humanized
<b>IgG type</b>	Human IgG1(K214R) - kappa
<b>Reactivity</b>	Human
<b>Target</b>	B7-H3
<b>Uniprot ID</b>	Q5ZPR3
<b>Description</b>	Anti-B7-H3(ifinatamab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only



## Anti-B7-H3(ifinatamab biosimilar) mAb ELISA

0.2  $\mu$ g of Human B7-H3, mFc-His tagged protein per well




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human B7-H3 Protein, mFc-His Tag (PME100012) can bind Anti-B7-H3(ifinatamab biosimilar) mAb (BME100532) in a linear range of 1.6-8.0 ng/mL.




Figure 2. B7-H3 protein is expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with 0.5  $\mu$ g/mL Anti-B7-H3(ifinatamab biosimilar) mAb (BME100532) (Blue histogram) or isotype control mAb (Red histogram) on HEK293 cells.

