

**PRODUCT INFORMATION**

<b>Common Name</b>	2857916, GSK2857916, J6M0, J6M0-mcMMAF, Unconjugated mAb
<b>Synonyms</b>	CD269;TNFRSF17;BCM;BCMA
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA; Flow Cyt
<b>Endotoxin</b>	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Humanized
<b>IgG type</b>	Human IgG1 - kappa
<b>Reactivity</b>	Human
<b>Target</b>	BCMA
<b>Uniprot ID</b>	Q02223
<b>Description</b>	Anti-BCMA (belantamab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals. Our unconjugated biosimilar monoclonal antibodies (mAbs) are based on the sequences outlined in relevant patents or scientific publications. These antibodies are in their native, unconjugated form, meaning they do not contain any payload or therapeutic agent attached. They are designed for use in research and development, and their performance has been tested as standalone molecules through comprehensive QC tests.
<b>Usage</b>	Research use only



## Anti-BCMA ( belantamab biosimilar) mAb ELISA




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human BCMA, mFc tagged protein PME100035 can bind Anti-BCMA Neutralizing antibody in a linear range of 0.64-80.0 ng/ml




Figure 2. Flow cytometry analysis with Anti-BCMA mAb 1  $\mu$ g/ml (belantamab) on K562 cells (Red histogram) or human BCMA positive K562 cells (Blue histogram). Human BCMA positive K562 cells were generated by human BCMA lentivirally transduced into K562 cell line and selected for monoclonal

