


**PRODUCT INFORMATION**

<b>Common Name</b>	JS004
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	BTLA;CD272
<b>Applications</b>	ELISA; Flow Cyt
<b>Endotoxin</b>	Less than 1.0 EU/ $\mu$ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Chimeric
<b>IgG type</b>	Human IgG4 - Kappa
<b>Reactivity</b>	Human
<b>Target</b>	BTLA
<b>Uniprot ID</b>	Q7Z6A9
<b>Description</b>	Anti-BTLA (icatolimab biosimilar)mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only




## Anti-BTLA (icatolimab biosimilar) mAb ELISA

0.1 µg of Human BTLA, His Tagged protein per well



**Figure 1.** ELISA plate pre-coated by 1 µg/ml (100 µl/well) Human BTLA , His tagged protein ([getskuurl sku="PME100524"]) can bind Anti-BTLA Neutralizing antibody (BME100054) in a linear range of 1-100 ng/ml.



**Figure 2.** Flow cytometry analysis with Anti-BTLA mAb 1 µg/ml (icatolimab) on HEK293 cells transfected with human BTLA (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram) .

