

PRODUCT INFORMATION

Clone ID DM92
Target BTN3A1

Synonyms BTN3A1; BTF5; CD277; BTN3.1; BT3.1

Host Species Rabbit

Description Anti-BTN3A1 antibody(DM92); Rabbit mAb

Delivery In Stock
Uniprot ID 000481
IgG type Rabbit IgG
Clonality Monoclonal
Reactivity Human

Applications ELISA; Flow Cyt

Recommended Dilutions

ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity

chromatography

Formulation & Reconstitution

Background

Conjugate

DIMA Disclaimer

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Unconjugated

The butyrophilin (BTN) genes are a group of major histocompatibility complex (MHC)-associated genes that encode type I membrane proteins with 2 extracellular immunoglobulin (Ig) domains and an intracellular B30.2 (PRYSPRY)

domain. Three subfamilies of human BTN genes are located in the MHC class I region: the single-copy BTN1A1 gene (MIM 601610) and the BTN2 (e.g.; BTN2A1; MIM 613590) and BTN3 (e.g.; BNT3A1) genes; which have undergone tandem duplication; resulting in 3 copies of each.

Usage Research use only

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under

patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

Email: info@dimabio.com Website: www.dimabio.com

actively scrutinizing all patent application to

ensure no IP infringement.








Figure 1. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human BTN3A1 protein, mFc-His tagged protein ([getskuurl sku="PME100056"]) can bind Rabbit anti-BTN3A1 monoclonal antibody (**clone: DM92**) in a linear range of 0.64-80 ng/ml.




Figure 2. Flow cytometry analysis with Anti-BTN3A1 (**DM92**) on HEK293 cells transfected with human BTN3A1 (**Red histogram**) or HEK293 transfected with irrelevant protein (**Blue histogram**).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

