

PRODUCT INFORMATION

Clone ID	3E7
Target	BTN3A3
Synonyms	BTF3
Host Species	Rabbit
Description	Anti-BTN3A3 antibody(3E7), Rabbit mAb
Delivery	In Stock
Uniprot ID	O00478
IgG type	Rabbit mAb
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1/100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The butyrophilin (BTN) genes are a group of major histocompatibility complex (MHC)-associated genes that encode type I membrane proteins with 2 extracellular immunoglobulin (Ig) domains and an intracellular B30.2 (PRYSPRY) domain. Three subfamilies of human BTN genes are located in the MHC class I region: the single-copy BTN1A1 gene (MIM 601610) and the BTN2 (e.g., BTN2A1; MIM 613590) and BTN3 (e.g., BNT3A3) genes, which have undergone tandem duplication, resulting in 3 copies of each (summary by Smith et al., 2010 [PubMed 20208008]).[supplied by OMIM, Nov 2010]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Flow cytometry analysis with 1 μ g/mL Anti-BTN3A3 (3E7) mAb on HEK293 cells transfected with human BTN3A3 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

