


## PRODUCT INFORMATION

<b>Common Name</b>	4A19, BMS-986340
<b>Synonyms</b>	CC-CKR-8;CCR-8;CDw198;CKRL1;CMKBR8;CMKBRL2;CY6;GPCRY6;TER1
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Humanized
<b>IgG type</b>	Human IgG1 - kappa
<b>Reactivity</b>	Human
<b>Target</b>	CCR8
<b>Uniprot ID</b>	P51685
<b>Description</b>	Anti-CCR8(BMS 986340) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only


### Anti-CCR8 (BMS 986340) mAb ELISA

0.2 µg of Human CCR8, mFc tagged protein per well



**Figure 1.** ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human CCR8 Protein, mFc Tag (PME101103) can bind Anti-CCR8 Neutralizing antibody (BME100115) in a linear range of 0.15–39.06 ng/mL.





**Figure 2.** Flow cytometry analysis with 1  $\mu$ g/mL Anti-CCR8 (BMS 986340) mAb (BME100115) on HEK293 cells transfected with Human CCR8 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

