


**PRODUCT INFORMATION**

<b>Common Name</b>	BT-062, BT062, nBT062-DM4
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	SDC1; Syndecan-1; CD138; SYND1; SDC
<b>Applications</b>	Flow Cyt
<b>Endotoxin</b>	Less than 1.0 EU/ $\mu$ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Chimeric
<b>IgG type</b>	Human IgG4 - Kappa
<b>Reactivity</b>	Human
<b>Target</b>	CD138
<b>Uniprot ID</b>	P18827
<b>Description</b>	Anti-CD138 (indatuximab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 $\mu$ m) prior to use.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only





**Figure 1.** HEK293 cell line transfected with irrelevant protein (**B**) and human CD138 (**A**) were surface stained with anti-CD138 neutralizing antibody 1 $\mu$ g/ml (indatuximab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.

