

PRODUCT INFORMATION

Clone ID	DM13
Target	CD22
Synonyms	SIGLEC-2; SIGLEC2
Host Species	Rabbit
Description	Anti-CD22 antibody(DM13); Rabbit mAb
Delivery	In Stock
Uniprot ID	P20273
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	CD22 (CD22 Molecule) is a Protein Coding gene. Diseases associated with CD22 include Refractory Hematologic Cancer and Hairy Cell Leukemia. Among its related pathways are Downstream signaling events of B Cell Receptor (BCR) and Hematopoietic cell lineage. Gene Ontology (GO) annotations related to this gene include carbohydrate binding. An important paralog of this gene is SIGLEC2.
Usage	Research use only






Figure 1. Expi 293 cell line transfected with irrelevant protein (**left**) and human CD22 (**right**) were surface stained with Rabbit anti- CD22 monoclonal antibody 1µg/ml (**clone: DM13**) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.





Figure 2. Flow cytometry data of serially titrated Rabbit anti-CD22 monoclonal antibody (**clone: DM13**) on Raji-cells. The Y-axis represents the mean fluorescence intensity (MFI), while the X-axis represents the concentration of IgG used.

Address: Wuhan Institute of Biotechnology B7, Biolake No.666
Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com






Figure 3. Affinity ranking of different Rabbit anti-CD22 mAb clones by titration of different concentration onto Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.




Figure 4. Phylogenetic analysis of amino acid sequence of different Rabbit Anti-CD22 mAb clones. **A)** Heavy chain and **B)** Light chain.

