

PRODUCT INFORMATION

Common Name	ACD22-VCMAE, DCDT2980S, FCU2803, RG-7593, RO5541072-000, Unconjugated mAb
Synonyms	CD22;SIGLEC2;BL-CAM;SIGLEC-2;Siglec2;SIGLEC2FLJ22814
Conjugate	Unconjugated
Applications	ELISA; Flow Cyt
Endotoxin	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	CD22
Uniprot ID	P20273
Description	Anti-CD22 (pinatuzumab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals. Our unconjugated biosimilar monoclonal antibodies (mAbs) are based on the sequences outlined in relevant patents or scientific publications. These antibodies are in their native, unconjugated form, meaning they do not contain any payload or therapeutic agent attached. They are designed for use in research and development, and their performance has been tested as standalone molecules through comprehensive QC tests.
Usage	Research use only



Anti-CD22 (pinatuzumab vedotin biosimilar) mAb ELISA




Figure 1. ELISA plate pre-coated by 2 μ g/ml (100 μ L/well) Human CD22, hFc-His tagged protein (PME100005) can bind Anti-CD22 Neutralizing antibody in a linear range of 3.2-400 ng/ml. In order to specifically detect BME100029, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. HEK293 cell line transfected with irrelevant protein (**B**) and human CD22 (**A**) were surface stained with Anti-CD22 mAb 1 μ g/ml (pinatuzumab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.

