


**PRODUCT INFORMATION**

<b>Common Name</b>	CFZ-533, NVP-CFZ533, OM11-62MF
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	CD40;Bp50;CDW40;MGC9013;TNFRSF5;p50
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Homo sapiens
<b>IgG type</b>	Human IgG1 - kappa
<b>Reactivity</b>	Human
<b>Target</b>	CD40
<b>Uniprot ID</b>	P25942
<b>Description</b>	Anti-CD40 (iscalimab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only





## Anti-CD40 (iscalimab biosimilar) mAb ELISA

0.2 µg of CD40, mFc-His Tagged protein per well



**Figure 1.** ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human CD40, mFc-His tagged protein ([getskuurl sku="PME100015"]]) can bind Anti-CD40 Neutralizing antibody in a linear range of 0.64-80.0 ng/ml.



**Figure 2.** HEK236 cell line transfected with irrelevant protein **(B)** and human CD40 protein **(A)** were surface stained with anti-CD40 neutralizing antibody 1µg/ml (iscalimab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.

