


## PRODUCT INFORMATION

<b>Clone ID</b>	DM164
<b>Target</b>	CD40 Ligand
<b>Synonyms</b>	CD40LG;CD154;CD40L;HIGM1;IGM;IMD3;T-BAM;TNFSF5;TRAP;gp39
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-CD40 Ligand antibody(DM164); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P29965
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Endotoxin</b>	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is expressed on the surface of T cells. It regulates B cell function by engaging CD40 on the B cell surface. A defect in this gene results in an inability to undergo immunoglobulin class switch and is associated with hyper-IgM syndrome.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr





**Figure 1.** Flow cytometry analysis with Anti-CD40L (DM164) on HEK293 cells transfected with human CD40L (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

