

PRODUCT INFORMATION

 Clone ID
 7E1

 Target
 CD45

Synonyms B220; CD45; CD45R; GP180; L-CA; LCA; LY5; T200

Host Species Rabbit

Description Anti-CD45 antibody(7E1), IgG1 Chimeric mAb

Delivery In Stock **Uniprot ID** P08575

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal
Reactivity Human
Applications IHC

Recommended Dilutions

IHC 1:100

Purification Purified from cell culture supernatant by affinity

chromatography

Formulation & Reconstitution

Background

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including

regulate a variety of cellular processes including cell growth; differentiation; mitosis; and oncogenic transformation. This PTP contains an extracellular domain; a single transmembrane

segment and two tandem intracytoplasmic catalytic domains; and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes; or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases; and thus functions as a regulator of cytokine

receptor signaling. Alternatively spliced transcripts variants of this gene; which encode distinct isoforms; have been reported.

Usage Research use only
Conjugate Unconjugated

njugate Unconjugated

All DIMA recombinant antibodies are genuinely

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

Email: info@dimabio.com

Website: www.dimabio.com

actively scrutinizing all patent application to

ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

DIMA Disclaimer








Figure 1. A. DMC100725 at $10\mu g/ml$ staining CD45 in Hut78 cells by IHC (SKU# DMC100725);B. DMC100725 at $10\mu g/ml$ staining CD45 in human tonsil tissue by IHC (SKU# DMC100725)




Figure 2. Flow cytometry analysis of antigen binding of anti-human CD45 mAb(DMC100725). (A) DMC100725 does not bind to CHO-S cells that do not express CD45. (B) A clear peak shift of DMC100725 was seen compared to the control when incubated with CD45-expressing THP-1 cells, indicating strong binding of DMC100725 to CD45. Antibodies were incubated at 5 μ g/mL.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

