Cat. No. BME100129



## **PRODUCT INFORMATION**

**Common Name** TJ-004309, TJC-4309, TJD-5

NT5E;5'-Nucleotidase;5'-NT;NT5;NTE **Synonyms** 

Conjugate Unconjugated **Applications** ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions** 

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

**Host Species** Chimeric Humanized IgG type Human IgG1 - kappa

Reactivity Human **Target** CD73 **Uniprot ID** P21589

**Description** Anti-CD73(uliledlimab biosimilar) mAb

**Delivery** In Stock

Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in

**Background** therapeutic or diagnostic procedures for humans

or animals.

Usage Research use only









## Anti-CD73 (uliledlimab biosimilar) mAb ELISA

0.2 µg of Human CD73, hFc tagged protein per well





Figure 1. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human CD73 Protein, hFc Tag (PME100829) can bind Anti-CD73 Neutralizing antibody (BME100129) in a linear range of 3.66–937.50 ng/mL. In order to specifically detect BME100129, mouse anti-human Fab-specific antibody was used as detection antibody.



**Figure 2.** Flow cytometry analysis with 15  $\mu$ g/mL Anti-CD73 (uliledlimab biosimilar) mAb (BME100129) on HEK293 cells transfected with Human CD73 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

