

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Common Name</b>                      | MEDI-9447, MEDI9447   |
| <b>Conjugate</b>                        | Unconjugated  |
| <b>Synonyms</b>                         | NT5E, 5'-Nucleotidase, 5'-NT, NT5, NTE  |
| <b>Applications</b>                     | ELISA, Flow Cyt   |
| <b>Endotoxin</b>                        | Less than 1.0 EU/ $\mu$ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.  |
| <b>Recommended Dilutions</b>            | ELISA 1:5000-10000, Flow Cyt 1:100  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.   |
| <b>Host Species</b>                     | Homo sapiens  |
| <b>IgG type</b>                         | Human IgG1(K214R,L234F,L235E,P331S)   |
| <b>Reactivity</b>                       | Human   |
| <b>Target</b>                           | CD73  |
| <b>Uniprot ID</b>                       | P21589  |
| <b>Description</b>                      | Anti-CD73(oleclumab biosimilar) mAb   |
| <b>Delivery</b>                         | In Stock  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature. |
| <b>Sterility</b>                        | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 $\mu$ m) prior to use.  |
| <b>Background</b>                       | Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.   |
| <b>Usage</b>                            | Research use only   |



## Anti-CD73(oleclumab biosimilar) mAb ELISA

0.2  $\mu$ g of Human CD73, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human CD73 Protein, hFc Tag (PME100829) can bind Anti-CD73(oleclumab biosimilar) mAb (BME100217) in a linear range of 0.13-16 ng/mL. In order to specifically detect BME100217, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis with 1 $\mu$ g/mL Anti-CD73(oleclumab biosimilar) mAb (BME100217) on HEK293 cells transfected with Human CD73 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

