

**PRODUCT INFORMATION**

<b>Clone ID</b>	353A3
<b>Target</b>	CLDN18.2,CLDN18.1
<b>Synonyms</b>	Claudin 18
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-CLDN18 antibody(353A3), IgG1 Chimeric mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P56856
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1/100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Endotoxin</b>	Less than 1.0 EU/ $\mu$ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Formulation &amp; Reconstitution</b>	Powder
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 $\mu$ m) prior to use.
<b>Background</b>	CLDN18 belongs to the large claudin family of proteins; which form tight junction strands in epithelial cells.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Flow cytometry analysis with 1 $\mu$ g/mL Anti-CLDN18 antibody(353A3) mAb on HEK293 cells transfected with human CLDN18.2 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis with 1 $\mu$ g/mL Anti-CLDN18 antibody(353A3) mAb on HEK293 cells transfected with human CLDN18.2 (Blue histogram) or HEK293 transfected with human CLDN18.1(Yellow histogram) or HEK293 transfected with irrelevant protein (Red histogram).

