

PRODUCT INFORMATION

Clone ID DM165 **Target** CLEC12A

Synonyms CLEC12A;MICL;CLL-1;CLL1;DCAL2;DCAL-2;CD371

Host Species Rabbit

Description Anti-CLEC12A antibody(DM165); Rabbit mAb

Delivery In Stock **Uniprot ID** Q5QGZ9 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human

Applications ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Reconstitution

Background

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the C-type lectin:C-type lectin-like domain (CTL:CTLD) superfamily. Members of this family share a common protein fold and have diverse functions;

such as cell adhesion; cell-cell signaling; glycoprotein turnover; and roles in inflammation and immune response. The protein encoded by this gene is a negative regulator of granulocyte and monocyte function. Several alternatively and processing variants of this gene have been spliced transcript variants of this gene have been

described; but the full-length nature of some of these variants has not been determined. This gene is closely linked to other CTL:CTLD superfamily members in the natural killer gene

complex region on chromosome 12p13.

Usage Research use only Conjugate Unconjugated


> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

> > Email: info@dimabio.com Website: www.dimabio.com

DIMA Disclaimer reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.



/+86-400-006-0995(China)








Figure 1. Flow cytometry analysis with Anti-CLEC12A (DM165) on HEK293 cells transfected with human CLEC12A (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

