Cat. No. DME100106



## **PRODUCT INFORMATION**

Clone ID DM106 **Target** CSF1R

**Synonyms** CSF1R;C-FMS;CD115;CSFR;FIM2;FMS;M-CSFR

**Host Species** Rabbit

Description Anti-CSF1R antibody(DM106); Rabbit mAb

**Delivery** In Stock **Uniprot ID** P07333 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human

**Applications** ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Formulation &

Storage & Shipping

**Background** 

Lyophilized from sterile PBS, pH 7.4. Normally 5 % 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature. The protein encoded by this gene is the receptor for colony stimulating factor 1; a cytokine which

controls the production; differentiation; and function of macrophages. This receptor mediates most if not all of the biological effects of this cytokine. Ligand binding activates the receptor kinase through a process of oligomerization and transphosphorylation. The encoded protein is a tyrosine kinase transmembrane receptor and member of the CSF1:PDGF receptor family of tyrosine-protein kinases. Mutations in this gene have been associated with a predisposition to myeloid malignancy. The first intron of this gene contains a transcriptionally inactive ribosomal protein L7 processed pseudogene oriented in the

opposite direction. Alternative splicing results in multiple transcript variants. Expression of a splice variant from an LTR promoter has been found in Hodgkin lymphoma (HL); HL cell lines and

> Email: info@dimabio.com Website: www.dimabio.com

anaplastic large cell lymphoma.

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)







Figure 1. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human CSF1R protein, His tagged protein ([getskuurl sku="PME100067"]) can bind Rabbit anti- CSF1R monoclonal antibody (clone: DM106) in a linear range of 0.6-60 ng/ml.



**Figure 2.** Flow cytometry analysis with Anti-CSF1R **(DM106)** on Expi293 cells transfected with human CSF1R(Blue histogram)or Expi293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

