

## **PRODUCT INFORMATION**

Clone ID 9A5
Target CSF1R

**Synonyms** CSF1R,C-FMS,CD115,CSFR,FIM2,FMS,M-CSFR

Host Species Rabbit

**Description** Anti-CSF1R antibody(9A5), IgG1 Chimeric mAb

Delivery In Stock Uniprot ID P07333

**IgG type** Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal
Reactivity Human
Applications Flow Cyt

Recommended Dilutions

Flow Cyt 1/100

Purification Purified from cell culture supernatant by affinity

chromatography

Formulation & Reconstitution

**Background** 

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is the receptor for colony stimulating factor 1, a cytokine which controls the production, differentiation, and function of macrophages. This receptor mediates most if not all of the biological effects of this cytokine. Ligand binding activates the receptor kinase through a process of oligomerization and transphosphorylation. The encoded protein is a

transphosphorylation. The encoded protein is a tyrosine kinase transmembrane receptor and member of the CSF1/PDGF receptor family of tyrosine-protein kinases. Mutations in this gene have been associated with a predisposition to myeloid malignancy. The first intron of this gene contains a transcriptionally inactive ribosomal protein L7 processed pseudogene oriented in the opposite direction. Alternative splicing results in multiple transcript variants. Expression of a splice variant from an LTR promoter has been found in Hodgkin lymphoma (HL), HL cell lines and

anaplastic large cell lymphoma.

Usage Research use only
Conjugate Unconjugated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under

patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

Email: info@dimabio.com Website: www.dimabio.com

actively scrutinizing all patent application to

ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

**DIMA Disclaimer** 








Figure 1. Flow cytometry analysis with  $1\mu g/mL$  Anti-CSF1R (9A5) mAb on HEK293 cells transfected with human CSF1R (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).



Email: info@dimabio.com Website: www.dimabio.com

