

PRODUCT INFORMATION

Clone ID **DMC443** CXCR7 **Target**

Atypical chemokine receptor 3;CXC-**Synonyms**

R7;CXCR-7;Chemokine orphan receptor 1;G-

protein coupled receptor 159;RDC-1

Host Species

Anti-CXCR7 antibody(DMC443); IgG1 Chimeric **Description**

mAb

In Stock Delivery P25106 **Uniprot ID**

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Background

Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the G-protein coupled receptor family. Although this protein was earlier thought to be a receptor for vasoactive intestinal peptide (VIP); it is now considered to be an orphan receptor; in that its endogenous ligand has not been identified. The

protein is also a coreceptor for human

immunodeficiency viruses (HIV). Translocations involving this gene and HMGA2 on chromosome 12 have been observed in lipomas.

Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)





Figure 1. Flow cytometry analysis with Anti-CXCR7 (DMC443) on Expi293 cells transfected with human CXCR7 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).



Email: info@dimabio.com Website: www.dimabio.com

