

PRODUCT INFORMATION

Common Name	AMG-757
Conjugate	Unconjugated
Synonyms	DLL3 and CD3
Applications	ELISA
Endotoxin	Less than 1.0 EU/ μ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	ELISA 1:5000-10000
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1(R292C,N297G,V302C)
Reactivity	Human
Target	DLL3 and CD3
Uniprot ID	Q9NYJ7 and P07766
Description	Anti-DLL3 and CD3(tarlatamab biosimilar) BsAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



Anti-DLL3 and CD3(tarlatamab biosimilar) BsAb ELISA

0.2 μ g of Human DLL3, His tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human DLL3 Protein, His Tag (PME100060) can bind Anti-DLL3 and CD3(tarlatamab biosimilar) BsAb (BME100264) in a linear range of 0.64-80 ng/mL.

