

**PRODUCT INFORMATION**

<b>Common Name</b>	AMG-757 without CD3
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	SCDO1
<b>Applications</b>	Flow Cyt
<b>Endotoxin</b>	Less than 1.0 EU/ $\mu$ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Humanized
<b>IgG type</b>	Human IgG1 - kappa
<b>Reactivity</b>	Human
<b>Target</b>	DLL3
<b>Uniprot ID</b>	Q9NYJ7
<b>Description</b>	Anti-DLL3(tarlatamab without CD3 biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only






Figure 1. Flow cytometry analysis with 15µg/mL Anti-DLL3(tarlatamab without CD3 biosimilar) mAb(BME100257) on 293T-DLL3 cells (Blue histogram) or isotype control mAb on 293T-DLL3 cells (Red histogram).

