

PRODUCT INFORMATION

Clone ID	OTI4C8
Target	DsRed-Monomer
Synonyms	N/A
Host Species	Mouse
Description	Carrier-free (BSA/glycerol-free) Mouse Monoclonal DsRed-Monomer Antibody, clone OTI4C8
Delivery	3-4week
Uniprot ID	N/A
IgG type	IgG1
Clonality	Monoclonal
Reactivity	N/A
Applications	WB IHC
Recommended Dilutions	WB 1:500, IHC 1:100
Purification	N/A
Endotoxin	Less than 1.0 EU/ μ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Formulation & Reconstitution	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μ m) prior to use.
Background	N/A
Usage	Research use only
Conjugate	Unconjugated




Figure 1. HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DsRed-Monomer (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 μ g per lane) were separated by SDS-PAGE and immunoblotted with anti-DsRed-Monomer.






Figure 2. Immunohistochemical staining of Frozen section Brain tissue from hGFAP-Cre, Rosa26 TdTomato mice using anti-DsRed-Monomer mouse monoclonal antibody (1:100). Secondary antibody: Alexa 488 goat anti-mouse.

DIMABIO CONFIDENTIAL

