

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | OTI3E12 |
| Target | E2-Crimson |
| Synonyms | N/A |
| Host Species | Mouse |
| Description | Carrier-free (BSA/glycerol-free) Mouse monoclonal E2-Crimson antibody, Clone OTI3E12 |
| Delivery | 3-4week |
| Uniprot ID | N/A |
| IgG type | IgG1 |
| Clonality | Monoclonal |
| Reactivity | N/A |
| Applications | WB |
| Recommended Dilutions | WB: 1:500 |
| Purification | N/A |
| Endotoxin | Less than 1.0 EU/ μ g by the LAL method. For <1 EU/mg requirements, please contact us for customization. |
| Formulation & Reconstitution | Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose) |
| Storage&Shipping | Store at -20°C. |
| Sterility | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μ m) prior to use. |
| Background | N/A |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY E2-Crimson (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-E2-Crimson (DME900020) 1:500.

DIMABIO CONFIDENTIAL

