

**PRODUCT INFORMATION**

<b>Clone ID</b>	OTI5A2 (formerly 5A2)
<b>Target</b>	EGFP
<b>Synonyms</b>	N/A
<b>Host Species</b>	Mouse
<b>Description</b>	Mouse monoclonal eGFP antibody, clone OTI5A2
<b>Delivery</b>	1-2week
<b>Uniprot ID</b>	N/A
<b>IgG type</b>	IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	N/A
<b>Applications</b>	WB
<b>Recommended Dilutions</b>	WB 1:500-1:2000
<b>Purification</b>	N/A
<b>Endotoxin</b>	Less than 1.0 EU/ $\mu$ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Formulation &amp; Reconstitution</b>	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
<b>Storage&amp;Shipping</b>	Store at -20°C.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 $\mu$ m) prior to use.
<b>Background</b>	N/A
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated




Figure 1. HEK293T cells were transfected with the pCMV6-ENTRY control (left lane) or pCMV6-ENTRY eGFP (right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5  $\mu$ g per lane) were separated by SDS-PAGE and immuno-blotted with 5A2 anti-eGFP monoclonal antibody (DME900006) (1:2000).



DIMABIO CONFIDENTIAL

