

**PRODUCT INFORMATION**

<b>Clone ID</b>	3G5
<b>Target</b>	Eribulin
<b>Synonyms</b>	N/A
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-Eribulin antibody(3G5); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	N/A
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	N/A
<b>Applications</b>	ELISA
<b>Recommended Dilutions</b>	ELISA 1/5000-10000
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Endotoxin</b>	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use.
<b>Background</b>	N/A
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr



### ELISA assay to evaluate Eribulin-Antibody 0.2 $\mu$ g Human IgG-Eribulin per well




Figure 1. Elisa plates were pre-coated with IgG-Eribulin (0.2 $\mu$ g/per well). Serial diluted anti-Eribulin monoclonal antibody (DME101050) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Eribulin monoclonal antibody binding with IgG-Eribulin is 1.012ng/ml.

