

PRODUCT INFORMATION

|                              |   |
|------------------------------|---|
| Clone ID                     | DMC460  |
| Target                       | FGFR4   |
| Synonyms                     | CD334; JTK2; TKF  |
| Host Species                 | Rabbit  |
| Description                  | Anti-FGFR4 antibody(DMC460); IgG1 Chimeric mAb  |
| Delivery                     | In Stock  |
| Uniprot ID                   | P22455  |
| IgG type                     | Rabbit/Human Fc chimeric IgG1   |
| Clonality                    | Monoclonal  |
| Reactivity                   | Human   |
| Applications                 | Flow Cyt  |
| Recommended Dilutions        | Flow Cyt 1:100  |
| Purification                 | Purified from cell culture supernatant by affinity chromatography   |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| Storage & Shipping           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| Background                   | The protein encoded by this gene is a tyrosine kinase and cell surface receptor for fibroblast growth factors. The encoded protein is involved in the regulation of several pathways; including cell proliferation; cell differentiation; cell migration; lipid metabolism; bile acid biosynthesis; vitamin D metabolism; glucose uptake; and phosphate homeostasis. This protein consists of an extracellular region; composed of three immunoglobulin-like domains; a single hydrophobic membrane-spanning segment; and a cytoplasmic tyrosine kinase domain. The extracellular portion interacts with fibroblast growth factors; setting in motion a cascade of downstream signals; ultimately influencing mitogenesis and differentiation. [provided by RefSeq; Aug 2017] |
| Usage                        | Research use only   |
| Conjugate                    | Unconjugated  |
| DIMA Disclaimer              | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.  |






Figure 1. Flow cytometry analysis with Anti-FGFR4 (DMC460) on HEK293 cells transfected with human FGFR4 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human FGFR4 mAb(DMC100460).  
(A) DMC100460 does not bind to SiHa cells that do not express FGFR4.  
(B) A clear peak shift of DMC100460 was seen compared to the control when incubated with FGFR4-expressing Huh7 cells, indicating strong binding of DMC100460 to FGFR4. Antibodies were incubated at 5  $\mu$ g/mL.

