

PRODUCT INFORMATION

Common Name	M2
Conjugate	Unconjugated
Synonyms	DDK□Flag□
Applications	ELISA, Flow Cyt
Endotoxin	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Mouse
IgG type	Mouse IgG1 - Kappa
Reactivity	N/A
Target	Flag
Uniprot ID	N/A
Description	Anti-Flag (BM301) Mouse mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).Lyophilized antibodies are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



Anti-Flag (BM301) Mouse mAb ELISA

0.2 μ g of Human LRP6(20-1370), Flag tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human LRP6(20-1370) Protein, Flag Tag (PME101719) can bind Anti-Flag (BM301) Mouse mAb (BME100093) in a linear range of 0.64-80 ng/mL.




Figure 2. Flow cytometry analysis with 0.5 μ g/mL Anti-Flag (BM301) Mouse mAb (BME100093) on HEK293 cells transfected with Flag-tagged protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

