

**PRODUCT INFORMATION**

<b>Clone ID</b>	1E11
<b>Target</b>	GFAP
<b>Synonyms</b>	ALXDRD
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-GFAP(7-66) antibody(1E11), Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P14136
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA
<b>Recommended Dilutions</b>	ELISA 1:5000-10000
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Endotoxin</b>	Less than 1.0 EU/ $\mu$ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 $\mu$ m) prior to use.
<b>Background</b>	This gene encodes one of the major intermediate filament proteins of mature astrocytes. It is used as a marker to distinguish astrocytes from other glial cells during development. Mutations in this gene cause Alexander disease, a rare disorder of astrocytes in the central nervous system. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Oct 2008]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr



### Anti-GFAP (1E11) mAb ELISA

0.1  $\mu\text{g}$  of Human GFAP (7-66) Protein, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1  $\mu\text{g/ml}$  (100  $\mu\text{l/well}$ ) Human GFAP(7-66) protein, hFc tagged protein (PME100666) can bind Rabbit anti-GFAP(7-66) monoclonal antibody(clone: 1E11) in a linear range of 1-50 ng/ml.

