

## **PRODUCT INFORMATION**

Clone ID DMC371 GPC3 **Target** 

DGSX; GTR2-2; MXR7; OCI-5; SDYS; SGB; SGBS; **Synonyms** 

SGBS1

**Host Species** Rabbit

**Description** Anti-GPC3 antibody(DMC371); IgG1 Chimeric mAb

**Delivery** In Stock **Uniprot ID** P51654

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Background

Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS)

contain a core protein anchored to the

cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26; and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi-

Behmel syndrome; also known as Simpson dysmorphia syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq; Sep 2009] References Fu Ying, Urban Daniel J, Nani Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting
Hepatocellular Carcinoma.[J] .Hepatology; 2019;
70: 563-576. Zhang Yi-Fan,Ho

Mitchell, Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular carcinoma.

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 1. Flow cytometry data of serially titrated anti-GPC3 monoclonal antibody (DMC371) on Huh7 cell line.

Email: info@dimabio.com Website: www.dimabio.com

