

PRODUCT INFORMATION

Clone ID **DMC500 Target HBEGF**

Synonyms DTR; DTS; DTSF; HEGFL

Host Species Rabbit

Anti-HBEGF antibody(DMC500); IgG1 Chimeric Description

mAb **Delivery** In Stock **Uniprot ID** Q99075

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Background

Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Growth factor that mediates its effects via EGFR; ERBB2 and ERBB4. Required for normal cardiac valve formation and normal heart function. Promotes smooth muscle cell proliferation. May be involved in macrophage-mediated cellular

proliferation. It is mitogenic for fibroblasts; but not endothelial cells. It is able to bind EGF receptor:EGFR with higher affinity than EGF itself and is a far more potent mitogen for smooth muscle cells than EGF. Also acts as a diphtheria toxin receptor.[UniProtKB:Swiss-Prot Function]

Usage Research use only

> Email: info@dimabio.com Website: www.dimabio.com










Figure 1. Flow cytometry analysis with Anti-HBEGF(DMC500) on Expi293 cells transfected with human HBEGF (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).





Figure 2. Flow cytometry analysis of antigen binding of anti-human HBEGF mAb(DMC100500). (A) DMC100500 does not bind to CHO-S cells that do not express HBEGF. (B) A clear peak shift of DMC100500 was seen compared to the control when incubated with HBEGF-expressing MDA-MB-231 cells, indicating strong binding of DMC100500 to HBEGF. Antibodies were incubated at 5 μ g/mL.

Email: info@dimabio.com Website: www.dimabio.com



/+86-400-006-0995(China)

