


PRODUCT INFORMATION


Common Name	JTX-2011
Conjugate	Unconjugated
Synonyms	ICOS;CD278;AILIM;Inducible T-cell costimulator
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	ICOS
Uniprot ID	Q9Y6W8
Description	Anti-ICOS (vopratelimab biosimilar) mAb
Delivery	In Stock
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



**Anti-ICOS (vopratelimab biosimilar) mAb ELISA**  
0.2 µg of Human ICOS, mFc-His Tagged protein per well



**Figure 1.** ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human ICOS, mFc-His tagged protein ([getskuurl sku="PME100026"]) can bind Anti-ICOS Neutralizing antibody in a linear range of 0.24-31.25ng/ml.



**Figure 2.** HEK293 cell line transfected with irrelevant protein (**B**) and human ICOS (**A**) were surface stained with anti-ICOS neutralizing antibody 1µg/ml (vopratelimab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.

