PRODUCT INFORMATION

Clone ID	DM183
Target	IL13RA1
Synonyms	CD213A1; CT19; IL-13Ra; NR4
Host Species	Rabbit
Description	Anti-IL13RA1 antibody(DM183); Rabbit mAb
Delivery	In Stock
Uniprot ID	P78552
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Endotoxin	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use.
Background	The protein encoded by this gene is a subunit of the interleukin 13 receptor. This subunit forms a receptor complex with IL4 receptor alpha; a subunit shared by IL13 and IL4 receptors. This subunit serves as a primary IL13-binding subunit of the IL13 receptor; and may also be a component of IL4 receptors. This protein has been shown to bind tyrosine kinase TYK2; and thus may mediate the signaling processes that lead to the activation of JAK1; STAT3 and STAT6 induced by IL13 and IL4.
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr



Figure 1. Flow cytometry analysis with Anti-IL13RA1 (DM183) on HEK293 cells transfected with human IL13RA1 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).