

**PRODUCT INFORMATION**

<b>Common Name</b>	HuMax-IL15, AMG 714, AMG-714, HuMax-IL-15, MAb146B7, PRV-015
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	IL-15
<b>Applications</b>	ELISA
<b>Endotoxin</b>	Less than 1.0 EU/ $\mu$ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Recommended Dilutions</b>	ELISA 1:5000-10000
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Homo sapiens
<b>IgG type</b>	Human IgG1(K214R,E356D,M358L) - Kappa
<b>Reactivity</b>	Human
<b>Target</b>	IL15
<b>Uniprot ID</b>	P40933
<b>Description</b>	Anti-IL15(ordesekimab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 $\mu$ m) prior to use.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only



### Anti-IL15(ordesekimab biosimilar) mAb ELISA

0.2 µg of Human IL15, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human IL15 Protein, hFc Tag (PME100882) can bind Anti-IL15(ordesekimab biosimilar) mAb (BME100253) in a linear range of 3.20-400 ng/mL. In order to specifically detect BME100253, mouse anti-human Fab-specific antibody was used as detection antibody.

### Anti-IL15(ordesekimab biosimilar) mAb ELISA

0.2 µg of Human IL15RA & IL15, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human IL15RA & IL15 Protein, hFc Tag (N120D) (PME101891) can bind Anti-IL15(ordesekimab biosimilar) mAb (BME100253) in a linear range of 1.6-8 ng/mL. In order to specifically detect BME100253, mouse anti-human Fab-specific antibody was used as detection antibody.



### Anti-IL15(ordesekimab biosimilar) mAb ELISA

0.2 µg of Human IL15RA and IL15, mFc tagged protein per well




Figure 3. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human IL15RA and IL15 Protein, mFc Tag (PME101926) can bind Anti-IL15(ordesekimab biosimilar) mAb (BME100253) in a linear range of 1.6-8.0 ng/mL.

