

PRODUCT INFORMATION

Common Name	ACZ885
Synonyms	IL-1 beta;IL1F2
Conjugate	Unconjugated
Applications	ELISA; Flow Cyt
Endotoxin	Less than 1.0 EU/ μ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Homo sapiens
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	IL1B
Uniprot ID	P01584
Description	Anti-IL1B(canakinumab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μ m) prior to use.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



Anti-IL1B (canakinumab biosimilar) mAb ELISA

0.1 μ g of Human IL1B, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/mL (100 μ L/well) Human IL1B Protein, hFc Tag (PME100838) can bind Anti-IL1B Neutralizing antibody (BME100116) in a linear range of 0.61-156.25 ng/mL. In order to specifically detect BME100116, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis under cell membrane permeable condition with 1 μ g/mL Anti-IL1B (canakinumab biosimilar) mAb (BME100116) on HEK293 cells transfected with Human IL1B protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

