

PRODUCT INFORMATION

Common Name	15D2, NN-8226, NNC-0109-0012
Synonyms	IL-20;Cytokine Zcyto10
Conjugate	Unconjugated
Applications	Flow Cyt
Endotoxin	Less than 1.0 EU/ μ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Homo sapiens
IgG type	Human IgG4 - Kappa
Reactivity	Human
Target	IL20
Uniprot ID	Q9NYY1
Description	Anti-IL20(fletikumab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μ m) prior to use.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only






Figure 1. Flow cytometry analysis under cell membrane permeable condition with 1 µg/mL Anti-IL20 (fletikumab biosimilar) mAb (BME100132) on HEK293 cells transfected with Human IL20 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

