

PRODUCT INFORMATION

Common Name 7F3com-2H2, MEDI-528

Synonyms Interleukin-9;IL-9;Cytokine P40;T-cell growth

factor P40

Conjugate Unconjugated

Applications ELISA; Flow Cyt

Recommended Dilutions

ELISA 1:5000-10000; Flow Cyt 1:100

Formulation & Reconstitution

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution.

Host Species Humanized

IgG type Human IgG1 – kappa

Reactivity Human
Target IL9

Uniprot ID P15248

Description Anti-IL9(enokizumab biosimilar) mAb

Delivery In Stock

Storage & Shipping

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Research grade biosimilar. Not for use in

Background therapeutic or diagnostic procedures for humans

or animals.

Usage Research use only

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

Email: info@dimabio.com Website: www.dimabio.com

patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to

ensure no IP infringement.





Anti-IL9 (enokizumab biosimilar) mAb ELISA

0.1 μg of Cynomolgus IL9, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/mL (100 μ L/well) Cynomolgus IL9 Protein, hFc Tag (PME-C100003) can bind Anti-IL9 Neutralizing antibody (BME100122) in a linear range of 2.44–156.25 ng/mL. In order to specifically detect BME100122, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis under cell membrane permeable condition with 1 μ g/mL Anti-IL9 (enokizumab biosimilar) mAb (BME100122) on Expi293 cells transfected with Human IL9 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com