

PRODUCT INFORMATION

Common Name	7F3com-2H2, MEDI-528
Synonyms	Interleukin-9;IL-9;Cytokine P40;T-cell growth factor P40
Conjugate	Unconjugated
Applications	ELISA; Flow Cyt
Endotoxin	Less than 1.0 EU/ μ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	IL9
Uniprot ID	P15248
Description	Anti-IL9(enokizumab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μ m) prior to use.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



Anti-IL9 (enokizumab biosimilar) mAb ELISA

0.1 μ g of Cynomolgus IL9, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/mL (100 μ L/well) Cynomolgus IL9 Protein, hFc Tag (PME-C100003) can bind Anti-IL9 Neutralizing antibody (BME100122) in a linear range of 2.44-156.25 ng/mL. In order to specifically detect BME100122, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis under cell membrane permeable condition with 1 μ g/mL Anti-IL9 (enokizumab biosimilar) mAb (BME100122) on HEK293 cells transfected with Human IL9 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

