

## **PRODUCT INFORMATION**

**Common Name** MCS-110, MCS110

**Synonyms** CSF-1;MCSF Conjugate Unconjugated **Applications** ELISA; Flow Cyt

Recommended

**Dilutions** 

ELISA 1:5000-10000; Flow Cyt 1:100

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution.

**Host Species** Humanized

IgG type Human IgG1 - kappa

Reactivity Human **Target** M-CSF **Uniprot ID** P09603

Anti-M-CSF(lacnotuzumab biosimilar) mAb **Description** 

**Delivery** In Stock

> Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in

**Background** therapeutic or diagnostic procedures for humans

or animals.

**Usage** Research use only

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

**DIMA Disclaimer** 

actively scrutinizing all patent application to ensure no IP infringement.


Email: info@dimabio.com Website: www.dimabio.com





## Anti-M-CSF (lacnotuzumab biosimilar) mAb ELISA

0.2 µg of Human M-CSF, His tagged protein per well



**Figure 1.** ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human M-CSF Protein, His Tag ([getskuurl sku="PME100569"]) can bind Anti-M-CSF Neutralizing antibody (**BME100104**) in a linear range of 2.44–156.25 ng/mL.

## Anti-M-CSF (lacnotuzumab biosimilar) mAb ELISA

0.2 µg of Human M-CSF, hFc tagged protein per well





Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human M-CSF Protein, hFc Tag (PME100550) can bind Anti-M-CSF Neutralizing antibody (BME100104) in a linear range of 9.77–156.25 ng/mL. In order to specifically detect BME100104, mouse anti-human Fab-specific antibody was used as detection antibody.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China) Email: info@dimabio.com Website: www.dimabio.com





**Figure 3.**Flow cytometry analysis with 1  $\mu$ g/mL Anti-M-CSF (lacnotuzumab biosimilar) mAb (BME100104) on HEK293 cells transfected with Human M-CSF protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

a

Email: info@dimabio.com Website: www.dimabio.com

