

PRODUCT INFORMATION

Common Name	OMS721
Conjugate	Unconjugated
Synonyms	MASP-2
Applications	Flow Cyt
Endotoxin	Less than 1.0 EU/ μ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Homo sapiens
IgG type	Human IgG4 - Lambda2
Reactivity	Human
Target	MASP2
Uniprot ID	O00187
Description	Anti-MASP2(narsoplimab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μ m) prior to use.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only






Figure 1. MASP2 protein is highly expressed on the surface of Jurkat cell membrane. Flow cytometry analysis with 15 µg/mL Anti-MASP2(narsoplimab biosimilar) mAb (BME100143) (Blue histogram) or isotype control mAb (Red histogram) on Jurkat cells.

DIMABIO CONFIDENTIAL

