

PRODUCT INFORMATION

Clone ID 7F12 MICA **Target**

MIC-A; PERB11.1 **Synonyms**

Host Species Rabbit

Description Anti-MICA antibody(7F12), IgG1 Chimeric mAb

Delivery In Stock **Uniprot ID** 029983

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human

Applications WB; Flow Cyt

Recommended

WB 1:1000; Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation &

8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes the highly polymorphic major histocompatability complex class I chain-related protein A. The protein product is expressed on the cell surface, although unlike canonical class I molecules it does not seem to associate with beta-2-microglobulin. It is a ligand for the NKG2-D type II integral membrane protein receptor. The protein functions as a stress-induced antigen that is broadly recognized by intestinal epithelial

Background gamma delta T cells. Variations in this gene have

been associated with susceptibility to psoriasis 1 and psoriatic arthritis, and the shedding of MICA-related antibodies and ligands is involved in the progression from monoclonal gammopathy of undetermined significance to multiple myeloma. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jan

20141

Usage Research use only Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under

> > Email: info@dimabio.com Website: www.dimabio.com

patent application. Any protein sequencing or **DIMA Disclaimer** reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)








Figure 1.Anti-MICA antibody (SKU# DMC100608) at 1/1000 dilution

Lane: HeLa (human cervical adenocarcinoma epithelial cell), whole cell lysate

Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

band size: 43 kDa





Figure 2. Flow cytometry analysis of antigen binding of anti-human MICA mAb(DMC100608).

(A) DMC100608 does bind to PC3 cells that weakly express MICA (B) A clear peak shift of DMC100608 was seen compared to the control when incubated with MICA-expressing Hela cells, indicating strong binding of DMC100608 to MICA. Antibodies were incubated at 10 ug/mL.



Email: info@dimabio.com Website: www.dimabio.com

