

PRODUCT INFORMATION

Clone ID	1G10
Target	MST1R
Synonyms	CD136;CDw136;NP6A3;PTK8;RON
Host Species	Rabbit
Description	Anti-MST1R antibody(1G10), Rabbit mAb
Delivery	In Stock
Uniprot ID	Q04912
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1/100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). This gene encodes a cell surface receptor for macrophage-stimulating protein (MSP) with tyrosine kinase activity. The mature form of this protein is a heterodimer of disulfide-linked alpha and beta subunits, generated by proteolytic cleavage of a single-chain precursor. The beta subunit undergoes tyrosine phosphorylation upon stimulation by MSP. This protein is expressed on the ciliated epithelia of the mucociliary transport apparatus of the lung, and together with MSP, thought to be involved in host defense. Alternative splicing generates multiple transcript variants encoding different isoforms that may undergo similar proteolytic processing. [provided by RefSeq, Jan 2016]
Background	
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Flow cytometry analysis with 2 μ g/mL Anti-MST1R (1G10) mAb on HEK293 cells transfected with human MST1R (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

