

## **PRODUCT INFORMATION**

**Common Name** IMC RON-8, IMC-RON8, RON8

CD136;CDw136;NPCA3;PTK8;RON **Synonyms** 

**Applications ELISA** 

Recommended

ELISA 1:5000-10000 **Dilutions** 

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

**Host Species** Homo sapiens

IgG type lgG1 Reactivity Human MST1R **Target Uniprot ID** Q04912

**Description** Anti-MST1R(narnatumab biosimilar) mAb

**Delivery** In Stock

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

Email: info@dimabio.com Website: www.dimabio.com

témperature.

Research grade biosimilar. Not for use in

**Background** therapeutic or diagnostic procedures for humans

or animals.

**Usage** Research use only





## Anti-MST1R (narnatumab biosimilar) mAb ELISA

0.2 µg of Human MST1R, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human MST1R Protein, hFc Tag (PME100821) can bind Anti-MST1R Neutralizing antibody (BME100118) in a linear range of 0.73–46.88 ng/mL. In order to specifically detect BME100118, mouse anti-human Fab-specific antibody was used as detection antibody.

Email: info@dimabio.com Website: www.dimabio.com

