

**PRODUCT INFORMATION**

<b>Common Name</b>	IMC RON-8, IMC-RON8, RON8
<b>Synonyms</b>	CD136;CDw136;NPCA3;PTK8;RON
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA, Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000, Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Homo sapiens
<b>IgG type</b>	Human IgG1 - kappa
<b>Reactivity</b>	Human
<b>Target</b>	MST1R
<b>Uniprot ID</b>	Q04912
<b>Description</b>	Anti-MST1R(narnatumab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only



## Anti-MST1R (narnatumab biosimilar) mAb ELISA

0.2 µg of Human MST1R, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human MST1R Protein, hFc Tag (PME100821) can bind Anti-MST1R Neutralizing antibody (BME100118) in a linear range of 0.73–46.88 ng/mL. In order to specifically detect BME100118, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis with 0.5 µg/mL Anti-MST1R(narnatumab biosimilar) mAb (BME100118) on HEK293 cells transfected with Human MST1R protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

